

AD-A037 531

PURDUE UNIV LAFAYETTE IND DEPT OF STATISTICS

F/G 12/1

MINIMAX ESTIMATION OF A MULTIVARIATE NORMAL MEAN WITH UNKNOWN C--ETC(U)

JUL 76 L J GLESER

AF-AFOSR-2350-72

UNCLASSIFIED

MIMEOGRAPH SER-460

AFOSR-TR-77-0177

NL

1 OF 1
AD
A037531




END

DATE
FILMED
4-77

3
B.S.

ADA037531



[Handwritten signature]


PURDUE UNIVERSITY

Approved for public release;
distribution unlimited.




DEPARTMENT OF STATISTICS

DIVISION OF MATHEMATICAL SCIENCES



AJ No.
DDC FILE COPY



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

Minimax Estimation of a Multivariate
Normal Mean with Unknown Covariance Matrix*

Leon Jay Gleser
Purdue University



Department of Statistics
Division of Mathematical Sciences
Mimeograph Series #460 ✓

July 1976

*Research supported in part by the Air Force Office of Scientific Research, Air Force Systems Command, USAF, under Grant AFOSR - 72-2350 ✓ at Purdue University.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
D C	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

291730

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR - TR - 77 - 0177	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) MINIMAX ESTIMATION OF A MULTIVARIATE NORMAL MEAN WITH UNKNOWN COVARIANCE MATRIX,		5. TYPE OF REPORT & PERIOD COVERED Interim rept.
7. AUTHOR(s) Leon Jay Gleser		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, IN 47907		8. CONTRACT OR GRANT NUMBER(s) AFOSR 72-2350
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 14p.		12. REPORT DATE Jul 76
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
14. Mimeograph Set - 460		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) +		
15. ✓ AF - AFOSR - 2350 - 72		
16. SUPPLEMENTARY NOTES (16) 2304 (17) A5		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) multivariate normal distribution, unknown covariance matrix, estimation of mean vector, quadratic loss, minimax estimator $(\delta - \theta)' Q (\delta - \theta)$		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let x be a p-variate ($p \geq 3$) vector, normally distributed with unknown mean θ and unknown covariance matrix Σ. Let $W: p \times p$ be distributed independently of x, and let W have a Wishart distribution with n degrees of freedom and parameter Σ. It is desired to estimate θ under the quadratic loss $(\delta - \theta)' Q (\delta - \theta)$ where Q is a known positive definite matrix. Under the condition that a lower bound for the smallest characteristic root of $Q \Sigma$ is known, a family of minimax estimators is developed.		

Use capital X
Sigma

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

291 730

Minimax Estimation of a Normal Mean

**Leon Jay Gleser
Department of Statistics
Mathematical Sciences Bldg.
Purdue University
West Lafayette, Indiana 47907**

Minimax Estimation of a Multivariate Normal Mean with Unknown Covariance Matrix

by

Leon Jay Gleser
Purdue University

ABSTRACT

Let x be a p -variate ($p \geq 3$) vector, normally distributed with unknown mean θ and unknown covariance matrix Σ . Let $W: p \times p$ be distributed independently of x , and let W have a Wishart distribution with n degrees of freedom and parameter Σ . It is desired to estimate θ under the quadratic loss $(\delta - \theta)'Q(\delta - \theta)$, where Q is a known positive definite matrix. Under the condition that a lower bound for the smallest characteristic root of $Q \Sigma$ is known, a family of minimax estimators is developed.

AMS 1970 Subject classification: Primary 62 C 99; secondary 62 F 10, 62 H 99.

Key words and phrases: Multivariate normal distribution, unknown covariance matrix, estimation of mean vector, quadratic loss, minimax estimator.

Minimax Estimation of a Multivariate Normal Mean with Unknown Covariance Matrix

by

Leon Jay Gleser
Purdue University

1. INTRODUCTION

Let $x: p \times 1$ be a normally distributed random vector with unknown mean θ and unknown covariance matrix Σ . Assume that we have an independent estimator $\hat{\Sigma} = n^{-1} W$ of Σ , where $W: p \times p$ has a Wishart distribution with n degrees of freedom and parameter $\Sigma = n^{-1} E(W)$. In the usual notation,

$$x \sim N(\theta, \Sigma), \quad W \sim \mathcal{W}_p(n, \Sigma). \quad (1)$$

We wish to estimate θ with an estimator $\delta(x, W)$ subject to the quadratic loss function

$$L(\delta, \theta, \Sigma) = (\delta - \theta)' Q (\delta - \theta) / \text{tr}(Q \Sigma) \quad (2)$$

Here, Q is a known $p \times p$ positive definite matrix, and $\text{tr}(A)$ denotes the trace of the matrix A . Note that $\text{tr}(Q \Sigma)$ is just a normalizing constant, chosen to give the estimator $\delta_0(x, W) = x$ constant risk. It is well known that δ_0 is a minimax estimator for this problem.

The limiting case of this problem where Σ is completely known (corresponding here to $n \rightarrow \infty$) has recently received a good deal of attention. [See Berger [1] for references.] The problem with Σ unknown and $Q = \Sigma^{-1}$ (which is not a special case for our problem because $Q = \Sigma^{-1}$ cannot be known) has also been studied by James and Stein [5], Lin and Tsai [6], Bock [2], and Efron and Morris [3,4], among others. However, the assumption that $Q = \Sigma^{-1}$ is rather artificial (it seems to be motivated only by invariance arguments), and does not seem to be of practical importance. A possibly more reasonable assumption to make relating Q and Σ is

that something is known about the characteristic roots of $Q\Sigma$. [Note that if $Q = \Sigma^{-1}$, all of the characteristic roots of $Q\Sigma$ are equal to 1.] In the present paper, we assume that there exists a known constant $K > 0$ such that

$$\text{ch}_p(Q\Sigma) \geq K, \text{ all } \Sigma > 0, \quad (3)$$

where

$$\text{ch}_1(A) \geq \text{ch}_2(A) \geq \dots \geq \text{ch}_p(A)$$

denote the ordered characteristic roots of the $p \times p$ symmetric matrix A .

We consider estimators of the form

$$\delta_h(x, W) = (I_p - h(x'W^{-1}x)Q^{-1}W^{-1})x, \quad (4)$$

where $h(u)$ is an absolutely continuous function on $[0, \infty)$. Our main result, which is proven in Section 2, is the following.

THEOREM 1. If (3) holds, then any estimator of the form (4) for which

(i) $u h(u)$ is nondecreasing in u ,

(ii) $0 \leq h(u) \leq 2(p-2)(n-p)K/(n-1)$, all $u \geq 0$,

dominates $\delta_0(x, W) = x$ in risk, and hence is minimax.

It is clearly of interest to determine what happens to estimators of the form (4) when the bound (3) can be violated. In Section 3 it is shown that when (3) does not hold, no estimator of the form (4) can be minimax. [Bock [2] has previously shown that for $Q = I_p$, no estimator of the form $h(x'W^{-1}x)x$ can be minimax.] It is conjectured that members of a certain family (see (36)) of estimators closely resembling the estimators (4) in form may be minimax, but no proof of this result is given.

2. PROOF OF THEOREM 1

Let

$$\Delta(\theta, \Sigma) = \text{tr}(Q\Sigma)E[L(\delta_h, \theta, \Sigma) - L(\delta_0, \theta, \Sigma)]. \quad (5)$$

Clearly if $\Delta(\theta, \Sigma) \leq 0$, all θ , all Σ satisfying (3), then δ_h is minimax for our problem.

Using the fact that $a'Qa - b'Qb = (a-b)'Q(a+b)$, the fact that $\delta_0(x, W) = x$, and (4), we obtain

$$\Delta(\theta, \Sigma) = E[h^2(x'W^{-1}x)x'W^{-1}Q^{-1}W^{-1}x] - 2E[h(x'W^{-1}x)x'W^{-1}(x-\theta)]. \quad (6)$$

Note that for any functions $g(x, W)$ for which $Eg(x, W)$ exists, we may write

$$E[g(x, W)] = E_W\{E_{X|W}[g(x, W)]\} = E_W\{E_X[g(x, W)]\}, \quad (7)$$

where $E_{X|W}[g(x, W)]$ denotes expectation over the conditional distribution of x given W , and E_W and E_X denote expectations over the marginal distributions of W and x respectively. The last equality in (7) holds since x and W are statistically independent. Further, using integration by parts term by term in the elements of x (with W treated as a fixed matrix), it can be shown (see Berger [1]) that

$$E_X[h(x'W^{-1}x)x'W^{-1}(x-\theta)] = E_X[h(x'W^{-1}x) \text{tr}W^{-1}] + 2E_X[h^{(1)}(x'W^{-1}x)x'W^{-1}\Sigma W^{-1}x], \quad (8)$$

where $h^{(1)}(u) = dh(u)/du$. [Note: We are assuming that $h(u)$ is differentiable; if not, a similar argument, using Riemann integration, produces a corresponding result; see Berger [1].]

From (6), (7), and (8), we have

$$\Delta(\theta, \Sigma) = E[h^2(x'W^{-1}x)x'W^{-1}Q^{-1}W^{-1}x] - 2h(x'W^{-1}x)\text{tr}W^{-1}\Sigma - 4h^{(1)}(x'W^{-1}x)x'W^{-1}\Sigma W^{-1}x]. \quad (9)$$

We now find a canonical representation for (9). Make the change of variables

$$y = \Sigma^{-1/2}x, \quad v = \Sigma^{-1/2}W\Sigma^{-1/2}, \quad (10)$$

where $\Sigma^{1/2}$ is any square root of Σ . Then

$$y \sim N(\eta, I_p), \quad V \sim \mathcal{W}_p(n, I_p), \quad (11)$$

where $\eta = \Sigma^{-1/2}\theta$. Further, y and V are statistically independent. From (9) and (10), with

$$Q^* = \Sigma^{1/2} Q \Sigma^{1/2},$$

and using arguments and notation analagous to that used to obtain (7), we have

$$\Delta(\theta, \Sigma) = E_y E_V [h^2 (y'V^{-1}y) y'V^{-1}(Q^*)^{-1}V^{-1}y - 2h(y'V^{-1}y)\text{tr}V^{-1} \quad (1)$$

$$- 4h^{(1)}(y'V^{-1}y)y'V^{-2}y] \quad (12)$$

Let Γ_y be $p \times p$ orthogonal with first row equal to $(y'y)^{-1/2}y'$. Let

$$U = \Gamma_y V \Gamma_y', \quad Q_y^* = \Gamma_y Q^* \Gamma_y'. \quad (13)$$

Then, given y , $U \sim \mathcal{W}_p(n, I_p)$, so that U and y are statistically independent. Partition U as

$$U = \begin{pmatrix} u_{11} & u_{21}' \\ u_{21} & U_{22} \end{pmatrix}, \quad u_{11}: 1 \times 1, \quad U_{22}: (p-1) \times (p-1),$$

and let

$$s = u_{11} - u_{21}' U_{22}^{-1} u_{21}, \quad t = U_{22}^{-1/2} \quad (14)$$

where $U_{22}^{1/2}$ is any square root of U_{22} . It is well known that s , t , and U_{22} are statistically independent, with

$$s \sim \chi_{n-p+1}^2, \quad t \sim N(0, I_{p-1}), \quad U_{22} \sim \mathcal{W}_{p-1}(n, I_{p-1}). \quad (15)$$

Further, $V^{-1} = \Gamma_y' U^{-1} \Gamma_y$ and

$$U^{-1} = s^{-1} \begin{pmatrix} 1 & -t' U_{22}^{-1/2} \\ -U_{22}^{-1/2} t & U_{22}^{-1/2} (s I_{p-1} + t t') U_{22}^{-1/2} \end{pmatrix}, \quad (16)$$

so that

$$y'V^{-1}y = s^{-1}y'y, \quad y'V^{-2}y = s^{-2}y'y(1 + t'U_{22}^{-1}t), \quad (17)$$

$$\text{tr}V^{-1} = \text{tr}U^{-1} = s^{-1} (1+t'U_{22}^{-1}t) + \text{tr}U_{22}^{-1}, \quad (18)$$

and

$$y'V^{-1}(Q^*)^{-1}V^{-1}y = s^{-2}y'y(1,-t'U_{22}^{-1/2})(Q_y^*)^{-1}(1,-t'U_{22}^{-1/2})'. \quad (19)$$

Under the distributional assumptions given in (15), it is known that $E(U_{22}^{-1}) = (n-p)^{-1}I_{p-1}$, so that

$$E\text{tr}U_{22}^{-1} = \text{tr}EU_{22}^{-1} = (n-p)^{-1}(p-1). \quad (20)$$

For any constant matrix A,

$$\begin{aligned} & E[(1,-t'U_{22}^{-1/2})A(1,-t'U_{22}^{-1/2})'] \\ &= E_{U_{22}} E_t \left\{ \text{tr} \left[A \begin{pmatrix} 1 & -t'U_{22}^{-1/2} \\ -U_{22}^{-1/2}t & U_{22}^{-1/2}t t' U_{22}^{-1/2} \end{pmatrix} \right] \right\} \\ &= E_{U_{22}} \text{tr} \left[A \begin{pmatrix} 1 & 0 \\ 0 & U_{22}^{-1} \end{pmatrix} \right] \\ &= \text{tr} \left[A \begin{pmatrix} 1 & 0 \\ 0 & (n-p)^{-1}I_{p-1} \end{pmatrix} \right] \end{aligned} \quad (21)$$

Taking $A = I_p$, the result (21) allows us to verify that

$$E(1 + t'U_{22}^{-1}t) = (n-p)^{-1}(n-1). \quad (22)$$

Taking $A = (Q_y^*)^{-1}$, the result (21) yields

$$\begin{aligned} & E[(1,-t'U_{22}^{-1/2})(Q_y^*)^{-1}(1,-t'U_{22}^{-1/2})'] \\ &= \text{tr}(Q_y^*)^{-1} \begin{pmatrix} 1 & 0 \\ 0 & (n-p)^{-1}I_{p-1} \end{pmatrix}. \end{aligned} \quad (23)$$

If in (12) we make the change of variables (13) and (14), and take account of the identities (17), (18), and (19), then by taking our expected values in the order $E_y E_s E_{t,U_{22}}$, and using (20), (22), and (23), we obtain

$$\begin{aligned}\Delta(\theta, \Sigma) = & (n-p)^{-1} E_y E_s [h^2 (s^{-1} y' y) s^{-2} y' y \tau(y, Q^*) \\ & - 2h(s^{-1} y' y) s^{-1} (n-1) - 2h(s^{-1} y' y) (p-1) \\ & - 4h^{(1)} (s^{-1} y' y) s^{-2} y' y (n-1)],\end{aligned}\quad (24)$$

where

$$\begin{aligned}\tau(y, Q^*) &= \text{tr}(Q_y^*)^{-1} \begin{pmatrix} n-p & 0 \\ 0 & I_{p-1} \end{pmatrix} \\ &= (n-p-1)(y' y)^{-1} y' (Q^*)^{-1} y + \text{tr}(Q^*)^{-1}.\end{aligned}\quad (25)$$

Finally, integrating by parts in s , we can show that

$$E_s h(s^{-1} y' y) = (n-p-1) E_s [s^{-1} h(s^{-1} y' y)] - 2 E_s [s^{-2} y' y h^{(1)}(s^{-1} y' y)], \quad (26)$$

which, when substituted in (24), yields the expression

$$\begin{aligned}\Delta(\theta, \Sigma) = & (n-p)^{-1} E_y E_s [h^2 (s^{-1} y' y) s^{-2} y' y \tau(y, Q^*) - 2p(n-p) s^{-1} h(s^{-1} y' y) \\ & - 4(n-p) h^{(1)} (s^{-1} y' y) s^{-2} y' y],\end{aligned}\quad (27)$$

where

$$y \sim N(n, I_p), \quad s \sim \chi^2_{n-p+1},$$

y and s are independent, $\eta = \Sigma^{-1/2} \theta$, $Q^* = \Sigma^{1/2} Q \Sigma^{1/2}$, and $\tau(y, Q^*)$ is given by (25). The expression (27) is the desired cononical form.

Now, we are ready to complete the proof of Theorem 1.

Let

$$r(u) = u h(u), \quad (28)$$

and note that

$$h^{(1)}(u) = \frac{r^{(1)}(u)}{u} - \frac{r(u)}{u^2}, \quad (29)$$

where $r^{(1)}(u) = dr(u)/du$. Substituting in (27), we obtain

$$\begin{aligned}\Delta(\theta, \Sigma) = & (n-p)^{-1} E_y \{ (y' y)^{-1} E_s [r^2 (s^{-1} y' y) \tau(y, Q^*) - 2(p-2)(n-p) r(s^{-1} y' y) \\ & - 4(n-p) s^{-1} r^{(1)}(s^{-1} y' y)] \} \\ \leq & (n-p)^{-1} E_y E_s \left[\frac{r(s^{-1} y' y)}{y' y} (\tau(y, Q^*) r(s^{-1} y' y) - 2(p-2)(n-p)) \right],\end{aligned}\quad (30)$$

since, by assumption (i) of Theorem 1, $r(u)$ is nondecreasing in u . Note from (3) and (25) that

$$\begin{aligned} \tau(y, Q^*) &\leq (n-1) \text{ch}_1[(Q^*)^{-1}] \leq (n-1) [\text{ch}_p(Q\Sigma)]^{-1} \\ &\leq (n-1) K^{-1}. \end{aligned} \quad (31)$$

Thus, applying assumption (ii) of Theorem 1, (30), and (31), we conclude that for all θ satisfying (3),

$$\Delta(\theta, \Sigma) \leq 0, \quad \text{all } \theta.$$

This completes the proof of Theorem 1.1.

We remark that our proof actually demonstrates the following.

THEOREM 2. Let an estimator $\delta_h(x, W)$ of the form (4) satisfy

(i) $u h(u)$ is nondecreasing in u ,

(ii) $0 \leq h(u) \leq 2(p-2)(n-p)Lu^{-1}$ all $u \geq 0$,

where $L > 0$ is a given constant. Then if Σ satisfies

$$\frac{(n-p-1)(\text{ch}_p(Q\Sigma))^{-1} + \text{tr}(Q\Sigma)^{-1}}{\leq L^{-1}}, \quad (32)$$

we have

$$\Delta(\theta, \Sigma) \leq 0, \quad \text{all } \theta,$$

and $\delta_h(x, W)$ is minimax.

Although Theorem 2 is more general than Theorem 1, the additional generality is unlikely to be of practical importance.

3. THE CASE WHERE Σ IS COMPLETELY UNRESTRICTED

When Σ is unrestricted, and (3) need not hold, then $\delta_0(x, W)$ is essentially the only estimator of the form (4) that can be minimax.

THEOREM 3. When Σ is unrestricted, no estimator of the form $\delta_h(x, W) = (I_p - h(x'W^{-1}x)Q^{-1}W^{-1})x$ can be minimax unless $h(u) = 0$ for almost all $u \geq 0$.

Proof. Note from (25) that

$$\tau(y, Q^*) \geq \text{tr}(Q^*)^{-1}, \quad \text{for all } y. \quad (33)$$

Now from (33) and (27),

$$\Delta(\theta, \Sigma) \geq \text{tr}(Q^*)^{-1} E[h^2(s^{-1}y'y)s^{-2}y'y] - 2(n-p)E[ps^{-1}h(s^{-1}y'y)-2h^{(1)}(s^{-1}y'y)s^{-2}y'y] \quad (34)$$

where the expected values in (34) are easily shown to depend only on $\theta'\Sigma^{-1}\theta$. Thus, if we choose a sequence $\{(\theta_i, \Sigma_i)\}$ of parameter values such that $\theta_i'\Sigma_i^{-1}\theta_i = c$, all i , and

$$\text{tr}(Q^*)^{-1} = \text{tr}(\Sigma_i)^{-1}Q^{-1} \rightarrow \infty, \text{ as } i \rightarrow \infty,$$

we see that unless

$$E[h^2(s^{-1}y'y)s^{-2}y'y] = 0, \text{ all } \theta'\Sigma^{-1}\theta = c, \quad (35)$$

we will have $\Delta(\theta_i, \Sigma_i) \rightarrow \infty$. Thus, for some parameter points $\Delta(\theta, \Sigma)$ will be positive (indeed, infinitely large), and hence $\delta_h(x, W)$ cannot be minimax. On the other hand, it is easy to show that (35) holds if and only if $h(u) = 0$ for almost all $u \geq 0$. This completes the proof.

Estimators of the form (4) do not perform well when any linear combination of the elements of x has low variability (implying that $ch_p(\Sigma)$ is small). To find a class of minimax estimators when Σ is unrestricted, we might think of modifying members of the class (4) to produce new estimators of the form

$$\delta_h^*(x, W) = (I_p - ch_p(n^{-1}QW)h(x'W^{-1}x)Q^{-1}W^{-1})x. \quad (36)$$

Assuming that $ch_p(n^{-1}QW)$ and $ch_p(Q\Sigma)$ are close in value (which should be true at least when n is large), any member of the class (36) will behave like the minimax estimator x when $ch_p(\Sigma)$ is small, and will behave like $\delta_{ch_p(Q\Sigma)h}$ otherwise. Thus, we have good intuitive reasons for conjecturing that a member of the class (36) of estimators is minimax provided that (i) $uh(u)$ is nondecreasing in u , and (ii) $0 \leq h(u) \leq 2(p-2)u$, all $u \geq 0$. Unfortunately, we have not yet been able to prove this conjecture. One can follow the steps used in Section 2, but unlike the result (24) obtained for the class (4), integration over t and U_{22} does not lead to any simplification. This lack of simplification is due to

the fact that $ch_p(n^{-1}QW)$, after the change of variables from (x, W) to (y, s, t, U_{22}) , is a complicated and nonlinear function of y, s, t , and U_{22} .

REFERENCES

- [1] BERGER, J. (1976). Minimax estimation of a multivariate normal mean under arbitrary quadratic loss. J. Multivariate Analysis 6
- [2] BOCK, M. E. (1975). Minimax estimators of the mean of a multivariate normal distribution. Ann. Statist. 3 209-218.
- [3] EFRON, B. AND MORRIS, C. (1976a). Families of minimax estimators of the mean of a multivariate normal distribution. Ann. Statist. 4 11-21.
- [4] EFRON, B. AND MORRIS, C. (1976b). Multivariate empirical Bayes and estimation of covariance matrices. Ann. Statist. 4 22-32.
- [5] JAMES, W. AND STEIN, C. (1960). Estimation with quadratic loss. Proc. 4th Berkeley Symp. Math. Stat. Prob. 1, 361-379. Univ. California Press, Berkeley.
- [6] LIN, PI-EHR AND TSAI, HUI-LIANG. (1973). Generalized Bayes minimax estimators of the multivariate normal mean with unknown covariance matrix. Ann. Statist. 1 142-145.